

UTTG-26-91

High Temperature Limit of the Confining Phase

Joseph Polchinski

Theory Group
Department of Physics
University of Texas
Austin, Texas 78712
bitnet:joe@utaphy

ABSTRACT

The deconfining transition in non-Abelian gauge theory is known to occur by a condensation of Wilson lines. By expanding around an appropriate Wilson line background, it is possible at large N to analytically continue the confining phase to arbitrarily high temperatures, reaching a weak coupling confinement regime. This is used to study the high temperature partition function of an $SU(N)$ electric flux tube. It is found that the partition function corresponds to that of a string theory with a number of world-sheet fields that diverges at short distance.

The large- N limit is one of the few analytic methods available in strongly coupled gauge theory, but has not yet proven solvable beyond two dimensions. Nevertheless, the topological nature of the perturbation expansion makes it tempting to believe that the large- N theory can be recast as a theory of noninteracting strings.¹ The principal tools that have been used to study this are the lattice strong coupling expansion and the loop equation. The strong coupling expansion can be written as a sum over noninteracting surfaces, but the complicated weights make the continuum interpretation difficult.² The loop equations have been argued to be equivalent to those for a string theory with additional world-sheet fields,³ but the mathematical intricacies of the loop equations make this claim difficult to evaluate. In this paper we will employ another tool, one which has proven useful in understanding the physics of both gauge theory and string theory. This is the study of the vacuum amplitude in a periodic Euclidean spacetime. We will focus on one compact dimension of period β . The vacuum amplitude can then be interpreted as the partition function at inverse temperature β .

The basic question we wish to ask is the number of degrees of freedom of a long flux tube at high energies. In recent work with Strominger,⁴ we have investigated this question at low energies. In this limit one expects an effective string description with $D - 2$ degrees of freedom, the transverse oscillations of the tube; here D is the spacetime dimension. In ref. 4, this was shown to be consistent with a covariant conformal field theory of central charge 26, as needed for a consistent quantization of the string. The theory of ref. 4 is an effective theory, nonrenormalizable and so invalid at sufficiently high energies. As one increases the energy, one would not be surprised to

find that the flux tube has additional degrees of freedom, the Liouville field⁵ and the elf field³ being possible candidates. The number of such fields at asymptotically high energies, determined from the partition function, would be an important piece of information. Further, since non-Abelian gauge theory is asymptotically free, one might hope that this could be obtained from a perturbative calculation.

There is a problem, pointed out by Thorn.⁶ The high temperature theory, in which one can calculate, is in a plasma phase and is separated by a phase transition from the low-temperature confining phase in which the flux tube exists. The point of this paper is that at large N it is possible to analytically continue the low temperature phase past the transition to arbitrarily high temperatures. In this way, one finds a regime in which confinement and asymptotic freedom are simultaneously present.

To begin, we recall a remarkable parallel between the statistical mechanics of string theory and of large- N gauge theory, as discussed in detail by Atick and Witten.⁷ In both theories the leading order free energy, g_{string}^{-2} or N^{-2} from the spherical topology, is β -independent. In string theory this occurs because the spherical Riemann surface cannot wind around the periodic dimension. In gauge theory it occurs because confinement restricts the spectrum to color singlets, whose number goes as N^0 . In both theories there is a transition at some β_c to a phase in which the leading order free energy is β -dependent.

In $SU(N)$ gauge theory, where we need not initially assume N to be large, the order parameters for this transition are the Wilson lines wrapping

k times around the compact dimension τ ,⁸

$$W_k(\vec{x}) = \frac{1}{N} \text{Tr } P \exp\left(i \int_0^{k\beta} d\tau A^\tau(\tau, \vec{x})\right). \quad (1)$$

The theory has a Z_N symmetry, consisting of gauge transformations g which are aperiodic by an element of the center,

$$g(\tau + \beta) = e^{2\pi i l/N} g(\tau). \quad (2)$$

These multiply the order parameters by phases $e^{2\pi i k l/N}$. At low temperature, the Wilson line two-point function falls exponentially,

$$W_k(\vec{R}) W_{-k}(0) \sim e^{-\beta T_k(\beta) R}, \quad R \rightarrow \infty. \quad (3)$$

The two-point function vanishes at infinity, so the expectation values of the order parameters are zero and the Z_N symmetry is unbroken. The Wilson line W_1 is an external quark source, while the W_k correspond to combinations of sources in higher representations. The $T_k(\beta)$ are thus the effective string tensions for flux in the various representations.

It is very useful to consider also a dual point of view, in which τ is regarded as a spatial coordinate. Then

$$M_k(\beta) = \beta T_k(\beta) \quad (4)$$

are the masses of winding states, states with electric flux in the periodic direction. At sufficiently small β these winding states become tachyonic (presumably the singly-wound M_1^2 becomes negative first) and the theory makes a transition to a phase in which the order parameters have expectation values and the Z_N symmetry is broken.^{8,9} These expectation values mean that an isolated quark source has finite energy: the theory no longer confines.

In the bosonic string theory there are also winding states, solitons with vertex operators $e^{ik\beta(\tau_R-\tau_L)/4\pi}$ and masses

$$m_k^2(\beta) = -8\pi T + k^2\beta^2 T^2, \quad (5)$$

with $T = T_1(\infty)$ being the string tension. These carry a $U(1)$ symmetry, with world-sheet current $(\partial_z\tau, -\partial_{\bar{z}}\tau)$.¹⁰ The phase structure is as in the gauge theory. At large β the m^2 are all positive, and the $U(1)$ symmetry is unbroken. At sufficiently small β , m_1^2 becomes negative and the theory makes a transition to a phase of broken $U(1)$ symmetry.^{11,7}

We are interested in the analytic continuation of the unbroken phase to small β . In string theory such a continuation is quite routine. Bosonic string perturbation theory is an expansion around an unstable vacuum, due to the $k = 0$ tachyon in eq. (5).¹² In formal studies of the perturbation theory this is usually an uninteresting artifact, and amplitudes are defined by analytic continuation from positive masses. This can be done for the $k \neq 0$ tachyons as well. In particular, the tree-level soliton masses-squared $m_k^2(\beta)$ given in eq. (5) are simply analytic in β .

Given the identical natures of the gauge and string transitions, the same continuation should be possible in gauge theory. We wish to expand the high-temperature gauge theory around a non-standard Wilson line background with unbroken Z_N (confinement), rather than the broken symmetry phase. Since the string continuation is carried out order by order in g_{string} , we will make the corresponding expansion, large- N , in the gauge theory.

To obtain an effective field theory of the Wilson lines, fix the gauge such

that the τ component of the vector potential is τ -independent and diagonal,

$$A_{ab}^\tau(\tau, \vec{x}) = \frac{\theta_a(\vec{x})}{\beta} \delta_{ab}, \quad (6)$$

and carry out the one-loop integral over the spacelike components of the vector potential. This leaves an effective three dimensional field theory of the $\theta_a(\vec{x})$,

$$\int [d\theta_a] e^{-S_{\text{eff}}}. \quad (7)$$

with action¹³

$$S_{\text{eff}} = \int d^3x \left\{ \frac{1}{2g^2\beta} \vec{\nabla}\theta_a \cdot \vec{\nabla}\theta_a + \frac{1}{24\pi^2\beta^3} \sum_{a,b=1}^N \theta_{ab}^2 (2\pi - \theta_{ab})^2 \right\}, \quad (8)$$

where θ_{ab} is $\theta_a - \theta_b$ modulo 2π , defined to lie between 0 and 2π . The gradient term is from the tree level action and the potential term is a one loop effect. In the regime we will consider, the symmetric phase at small g , this is an honest low energy effective theory: the θ_a have masses of order $(g^2N)^{1/2}\beta^{-1}$, while the spacelike gauge bosons which have been integrated out have masses of order β^{-1} .¹⁴ There are corrections of higher order in g^2 , and also corrections of higher order in $\beta^2\nabla^2$. Since the one loop term arises from the scale β , the coupling in the action should be understood as $g(\beta)$.

In the large- N limit, the θ_a become infinite in number and it becomes appropriate to consider the normalized density¹⁵

$$\rho(\theta, \vec{x}) = \frac{1}{N} \sum_a \delta(\theta - \theta_a(\vec{x})). \quad (9)$$

The action becomes

$$\begin{aligned} S_c = & N^2 \int d^3x \int_0^{2\pi} d\theta \frac{1}{2g^2N\beta\rho} (\partial_\theta^{-1} \vec{\nabla}\rho)^2 \\ & + \frac{N^2}{24\pi^2\beta^3} \int d^3x \int \int_0^{2\pi} d\theta_1 d\theta_2 \rho(\theta_1) \rho(\theta_2) \theta_{12}^2 (2\pi - \theta_{12})^2. \end{aligned} \quad (10)$$

As usual in the large- N limit, this is of order N^2 with g^2N held fixed. The potential favors all eigenvalues being equal,

$$\rho_b(\theta, \vec{x}) = \delta(\theta - \theta_0), \quad (11)$$

breaking the $U(1)$ of $\theta \rightarrow \theta + \epsilon$. The symmetric phase is completely determined by the requirement that the eigenvalue distribution be $U(1)$ -invariant,

$$\rho_s(\theta, \vec{x}) = \frac{1}{2\pi}. \quad (12)$$

Expanding around the symmetric point,

$$\rho(\theta, \vec{x}) = \frac{1}{2\pi} \left\{ 1 + \sum_{\substack{k=-\infty \\ k \neq 0}}^{\infty} \rho_k(\vec{x}) e^{-ik\theta} \right\}, \quad (13)$$

the quadratic part of the action becomes

$$S_c^{(2)} = N^2 \sum_{\substack{k=-\infty \\ k \neq 0}}^{\infty} \int d^3x \left\{ \frac{1}{2g^2N\beta k^2} (\vec{\nabla} \rho_k)^2 - \frac{1}{\pi^2 \beta^3 k^4} \rho_k^2 \right\}. \quad (14)$$

The Wilson line is simply $W_k(\vec{x}) = \rho_k(\vec{x})$ so we can read off the winding state masses

$$M_k^2 = -\frac{2g^2N}{\pi^2 \beta^2 k^2}. \quad (15)$$

This pattern is quite a bit different from that of the critical string, eq. (5). As a check, both spectra depend on the parameters k and β in the combination $k\beta$.¹⁶ This is because a *planar* surface winding k times around a dimension of period β is the same as one winding once around a dimension of period $k\beta$. We can therefore restrict attention to $k = 1$.

It is convenient that the quantity the we are interested in, the number of degrees of freedom of a long $k = 1$ string, is directly related by a modular

transformation to $M_1^2(\beta)$.¹⁷ Suppose that the density of states of the flux tube is equal to that of n free world-sheet scalars, so the spectrum of a tube with ends fixed at 0 and $\vec{R}$ is

$$E^2 = T^2 R^2 + 2\pi T \sum_{a=1}^n \left(-\frac{1}{24} + \sum_{m=1}^{\infty} m N_{am} \right), \quad (16)$$

where T is the string tension, a labels the n scalars, m labels the harmonic, and N_{am} is the occupation number. We have included the Casimir constant $-\frac{1}{24}$ for good form, although it does not matter much at high temperature. The partition function $\text{Tr}(e^{-\beta E})$ can be found by standard methods.¹⁸ It is of the form (3) with

$$\beta^2 T_1^2(\beta) \equiv M_1^2(\beta) = -\frac{n\pi T}{3} + \beta^2 T^2. \quad (17)$$

At sufficiently small β this is always negative, the entropy of fluctuations overwhelming the string tension and favoring long strings. This is the usual picture of the Hagedorn transition, equivalent to the symmetry-breaking picture presented earlier.^{11,7}

Comparing equations (15) and (17), we see that the partition function of the flux tube is not consistent with any fixed number of degrees of freedom. Rather, as we go to smaller β we excite more and more degrees of freedom, with

$$n_{\text{eff}}(\beta) \sim \frac{6g^2(\beta)N}{\pi^3 T \beta^2}. \quad (18)$$

This is our main result. Any string theory equivalent to large- N gauge theory will have a number of fields which diverges at short distances.¹⁹ In particular, various proposals that the gauge theory might be equivalent to a string theory with a finite number of world-sheet fields would seem to be ruled out. This

will extend to the spectra of mesons and glueballs as well: the number of states at high energy will greatly exceed that in a string theory.

This is complementary to the result obtained by Atick and Witten in a very similar way.⁷ They show that at high energy string theory has fewer degrees of freedom than any field theory. It should then be no surprise that the large- N gauge theory, which is a field theory with moreover an infinite number of fields, should correspond to a string theory with an unusually large number of degrees of freedom. It is remarkable that we have been able to use perturbation theory to obtain the number of degrees of freedom of an object which exists only nonperturbatively. We have little physical intuition for how this is possible, but experience with string theory gives us confidence that the approach is correct. The high temperature continuation of the confining phase is quite similar to the Eguchi-Kawai reduction of the lattice theory,²⁰ as we will discuss further below.

Now let us make some general remarks about the program to reformulate the large- N theory as a string theory. There are at least three ways in which the flux tube might be unlike a fundamental string. First, it might have contact interactions, such as self-avoidance, which would be nonlocal on the world-sheet. The large- N factorization property²¹ shows that this is not the case, since factorization holds even for sources in close proximity. This also follows at least formally from the strong-coupling sum over surfaces representation.²

Second, the world-sheet might have holes: the large- N perturbation theory looks like a net rather than a surface. Here again the large- N flux tube is indistinguishable from a fundamental string:⁷ as already noted the spherical

amplitude in a spacetime of large period β is β -independent in both theories, indicating no holes, and in both theories holes develop spontaneously at small β from the condensation of winding states.

Third, the flux tube might be fat, unlike the infinitely thin fundamental string. Indeed, the number of degrees of freedom (18) grows as β^{-2} , aside from the slow running of the coupling. This is just what would be seen if the flux tube were a *three*-dimensional object with a thickness of order $T^{-1/2}$. Nevertheless, we believe that these degrees of freedom are actually internal, and that the large- N string has in some sense a zero intrinsic thickness. The first argument for this is the formal lattice representation as a sum over surfaces.² The second comes from another application of the high-temperature continuation. Consider a large Wilson loop lying in a plane, and compactify the two dimensions orthogonal to the plane. If the flux tube had an intrinsic thickness, we would expect that reducing the periods in the orthogonal directions would squeeze the tube and increase its energy, reducing the Wilson loop expectation value. In fact, the expectation value is independent of the size of the orthogonal directions at large N , because a planar surface cannot wind around these dimensions in the symmetric phase.²²

We have used the high temperature continuation to count the number of degrees of freedom in the flux tube and to argue that the tube has zero thickness in the large- N limit. This should be equivalent to a *two-dimensional* field theory with an infinite number of fields. In fact, the high temperature continuation makes it possible to construct this theory: compactify the orthogonal dimensions to very small radii while remaining in the symmetric phase. The resulting theory is essentially the dimensional reduction of the

gauge theory to two dimensions. The degrees of freedom are the N^2 gauge fields $A^\mu(\tau, \sigma)$; the main subtlety seems to be in the expansion around the symmetric point. Going further, compactifying all but one dimension yields a matrix quantum mechanics of $A^\mu(\tau)$ whose spectrum is equivalent to the original gauge theory. This result is already known from the Eguchi-Kawai reduction,^{20,23} which we see can be interpreted as the analytic continuation of the confining phase. This reduction has not yet proven analytically tractable, but perhaps a reexamination is in order.

Acknowledgements

I would like to thank L. McLerran, D. Minic, A. Strominger, and Z. Yang for helpful discussions. Much of this work was carried out while the author was a guest at the Aspen Center for Physics. This research was supported in part by the Robert A. Welch Foundation and NSF Grant PHY 9009850.

References

1. G. 't Hooft, Nucl. Phys. **B72**, 461 (1974); **B75**, 461 (1974).
2. V. A. Kazakov, Phys. Lett. **B128**, 316 (1983); V. I. Kostov, Phys. Lett. **B138**, 191 (1984); K. H. O'Brien and J.-B. Zuber, Nucl. Phys. **B253**, 621 (1985).
3. For a review see A. A. Migdal, Phys. Rep. **102**, 199 (1983).
4. J. Polchinski and A. Strominger, "Effective String Theory," Texas preprint UTTG-17-91 (1991).
5. A. M. Polyakov, Phys. Lett. **B103**, 207 (1981).
6. C. Thorn, Phys. Lett. **B99**, 458 (1981); R. D. Pisarski, Phys. Rev. **D29**, 1222 (1984).
7. J. Atick and E. Witten, Nucl. Phys. **B310**, 291 (1981).
8. A. M. Polyakov, Phys. Lett. **B72**, 447 (1978); L. Susskind, Phys. Rev. **D20**, 2610 (1979); for a review see B. Svetitsky, Phys. Rep. **132**, 1 (1986).
9. It is of some interest whether the transition is first order, occurring before $T_1(\beta)$ actually reaches zero, but this will not be relevant for us.
10. This is the $U(1)$ which couples to winding number. Both the gauge theory and the string theory have the additional $U(1)$ symmetry of τ -translations. This is unbroken in both phases.

11. B Sathiapalan, Phys. Rev. **D35**, 3277 (1987); Ya. I. Kogan, JETP Lett. **45**, 709 (1987); K. H. O'Brien and C.-I. Tan, Phys. Rev. **D36**, 1184 (1987).
12. The $k = 0$ tachyon is neutral under the $U(1)$. It is not clear whether it has any analog in the gauge theory.
13. D. J. Gross, R. D. Pisarski, and L. G. Yaffe, Rev. Mod. Phys. **53**, 43 (1981). See also N. Weiss, Phys. Rev. **D24**, 475 (1981); **D25**, 2667 (1982).
14. This is in contrast to the unconfined plasma phase, where the spacetime vector potential has a magnetic mass $g^2 N \beta^{-1}$.
15. E. Brézin, C. Itzykson, G. Parisi, and J.-B. Zuber, Comm. Math. Phys. **59**, 35 (1978); A. Jevicki and B. Sakita, Nucl. Phys. **B165**, 511 (1980).
16. In particular, a careful treatment of the loop corrections would show that the coupling g in eq. (15) must be evaluated at the scale $k\beta$.
17. This sort of relation between ground state energy and asymptotic degrees of freedom is well-known in string theory. See, for example, J. Cardy, Nucl. Phys. **B270** [FS16], 186 (1986). For a recent discussion, see D. Kutasov and N. Seiberg, Nucl. Phys. **B358**, 600 (1991).
18. R. Hagedorn, Nuovo Cim. Suppl. **3**, 147 (1965). To derive eq. (17), one can use the integral

$$e^{\beta E} = \frac{\beta}{2\pi} \int_0^\infty \frac{ds}{s^{3/2}} e^{(E^2 s + \beta^2 s^{-1})/2}$$

to remove the square root from the exponent, carry out the sum over occupation numbers, use the asymptotic formula

$$\prod_{m=1}^{\infty} (1 - x^m) \stackrel{x \rightarrow 1^-}{\sim} e^{-\pi^2/6 \ln x},$$

and then use the above integral again.

19. It may be unfamiliar that the number of degrees of freedom can vary with world-sheet scale, since the world-sheet theory is conformally invariant. The point is that in the noncritical string the conformal invariance is broken by the expectation value of the metric, e^ϕ or $\partial_z X^\mu \partial_{\bar{z}} X_\mu$, making massive fields possible.
20. T. Eguchi and H. Kawai, Phys. Rev. Lett. **48**, 1063 (1982). See ref. 3 for a review.
21. E. Witten, in NATO Advanced Studies Institute Series—Series B: Physics, Vol. 59, eds. G. 't Hooft, et. al. (Plenum, 1979).
22. This property of the confining phase was derived from the lattice Schwinger-Dyson equations by A. Gocksch and F. Neri, Phys. Rev. Lett. **B50**, 1099 (1983).
23. A continuum version of the Eguchi-Kawai reduction was discussed in D. J. Gross and Y. Kitazawa, Nucl. Phys. **B206**, 440 (1982) and in A. Migdal, Phys. Lett. **B116**, 425 (1982).